
The Case for Postpartum Warming Foods

How traditional food cultures worldwide converge on the same postpartum wisdom—and what modern science says about why

By *Monika Sudakov*

Introduction

In China, a new mother enters *zuo yue zi*—"sitting the month"—and is fed warming soups, ginger-infused broths, and slow-cooked pork knuckle with vinegar and sesame oil. In Korea, she drinks *miyeokguk*, a seaweed soup believed to purify the blood and promote milk production. In India, Ayurvedic tradition prescribes *ghee*-laden foods, warm spiced milk, and easily digestible grains. In Mexico, the *cuarentena*—the forty days—centers on *caldos* (broths), *atoles* (warm corn-based drinks), and gentle spiced stews.¹

These traditions developed independently across thousands of years, on different continents, in cultures with no contact with one another. Yet they converge on the same fundamental principles: warm food. Slow-cooked food. Nutrient-dense food. Food that is soft, easy to digest, and delivered by others. The consistency of this convergence is not coincidence. It is evidence.

This paper maps traditional postpartum food practices from around the world to what modern nutritional science, gastroenterology, and immunology now understand about the postpartum body. The conclusion is striking: these traditions got it right, and modern Western postpartum culture—which offers no structured nutritional guidance and expects new mothers to feed themselves—has it wrong.

A Global Survey of Postpartum Food Traditions

China: Zuo Yue Zi (坐月)

The earliest recorded references to *zuo yue zi* appear in classical Chinese medical texts from the Han Dynasty (206 BCE–220 CE). The practice rests on the Traditional Chinese Medicine concept that childbirth depletes the body's *qi* (vital energy) and blood, leaving the mother in a "cold" state that must be corrected through warming interventions.²

The postpartum diet centers on warming foods: pigs' feet and peanut soup (believed to promote lactation), ginger-fried rice, sesame oil chicken, red date tea, and rich bone broths simmered for hours. Cold foods—raw vegetables, cold drinks, fruits served at room temperature—are strictly avoided. The foods are classified not by temperature alone but by their *thermal nature* in TCM: their effect on the body's internal heat balance.³

Research published in *BMC Pregnancy and Childbirth* found that *zuo yue zi* remains widely practiced among Chinese women, both in China and in diaspora communities worldwide. A qualitative study of Chinese mothers in Switzerland found that all participants were aware of the practice, and a majority continued it to varying extents, adapting traditional practices to available ingredients and circumstances.⁴

Korea: Miyeokguk and the Seaweed Tradition

In Korea, *miyeokguk*—seaweed soup—is so strongly associated with birth that it is also served on birthdays as a reminder of one's mother's sacrifice. The soup is made with *miyeok* (wakame seaweed), beef broth, garlic, and sesame oil. It is eaten daily for the first several weeks postpartum, sometimes for all three meals.

The tradition has a nutritional logic that modern science validates: wakame seaweed is rich in iodine (essential for thyroid function, which is stressed postpartum), calcium (depleted by pregnancy and lactation), iron (lost through birth), and fucoidan (a sulfated polysaccharide with anti-inflammatory properties). The warm broth delivery system supports hydration and mineral absorption.⁵

India: Ayurvedic Postpartum Nutrition

Ayurvedic tradition classifies the postpartum period as a *vata* state—characterized by dryness, coldness, and instability. The prescribed antidote mirrors the Chinese approach: warm, oily, grounding foods. Ghee features prominently, both for its caloric density and for the Ayurvedic belief that it lubricates tissues and supports *ojas* (vitality). Warm spiced milk with turmeric and saffron, soft-cooked rice with lentils (*khichdi*), and ginger-infused preparations are staples.⁶

The Ayurvedic emphasis on digestibility is particularly noteworthy. Traditional texts specify that postpartum foods should be cooked until very soft, often reduced to semi-liquid consistency, to minimize the digestive burden on a depleted body. This aligns precisely with modern gastroenterological understanding of postpartum gut function.

Latin America: La Cuarentena

In many Latin American traditions, the *cuarentena*—the forty days—is a recognized period of rest and community care. The postpartum diet centers on *caldos* (broths, especially chicken), *atole* (warm corn-based drinks), and gentle stews. Cold foods and drinks are avoided. The preparation and delivery of these meals is a communal responsibility, typically managed by the mother's own mother, sisters, and neighbors.

The communal meal delivery component is as significant as the nutritional content. The *cuarentena* is not merely a dietary prescription—it is a social infrastructure that ensures the new mother is never required to feed herself during the recovery period.

"Nearly every culture that developed a postpartum food tradition independently arrived at the same conclusion: warm, slow-cooked, nutrient-dense food, delivered by others. When diverse civilizations converge on the same answer, it is worth asking what question they were all responding to."

— Monika Sudakov, Founder, Mothership

What Modern Science Says About Warming Foods

Gut Permeability and the Postpartum Microbiome

Pregnancy alters the gut microbiome in ways that persist into the postpartum period. Research has shown increased intestinal permeability ("leaky gut") during late pregnancy and early postpartum, likely mediated by progesterone withdrawal and inflammatory cytokine shifts. This increased permeability means the postpartum gut is more vulnerable to irritation from difficult-to-digest foods and more responsive to soothing, anti-inflammatory inputs.⁷

Warm, slow-cooked foods directly address this vulnerability. Prolonged cooking breaks down complex proteins and fibers into more bioavailable forms, reducing the digestive burden. Bone broth, a centerpiece of virtually every traditional postpartum diet, provides gelatin (cooked collagen) and amino acids like glutamine and glycine that research associates with intestinal barrier support. Glutamine serves as the primary fuel source for enterocytes (intestinal lining cells), while glycine has demonstrated anti-inflammatory properties in both animal and human studies.⁸

Inflammation and the Thermal Hypothesis

The TCM classification of postpartum as a "cold" state finds an intriguing parallel in modern immunology. Childbirth triggers a significant inflammatory cascade—the body's response to the physical trauma of delivery and the rapid hormonal reorganization that follows. C-reactive protein, IL-6, and other inflammatory markers are elevated in the early postpartum period.⁹

Many of the warming ingredients featured in traditional postpartum diets have documented anti-inflammatory properties. Ginger contains gingerols and shogaols, which inhibit prostaglandin synthesis and NF-κB activation. Turmeric's curcumin is one of the most studied natural anti-inflammatory compounds in the pharmacological literature. Garlic's allicin modulates multiple inflammatory pathways. Sesame oil contains sesamin, which has demonstrated anti-inflammatory and antioxidant effects.¹⁰

These are not folk remedies grasping at plausibility. These are bioactive compounds with well-characterized mechanisms of action, consumed in the forms (cooked, dissolved in broth, combined with fats that enhance bioavailability) that traditional food wisdom prescribed centuries before the mechanisms were understood.

Nutrient Bioavailability and Cooking Method

Slow cooking does more than make food tender. It fundamentally alters nutrient bioavailability. Prolonged heating breaks down plant cell walls, releasing bound minerals. It denatures proteins into more digestible forms. It dissolves collagen from bones into gelatin, converting an insoluble structural protein into a water-soluble, gut-accessible form.¹¹

For postpartum mothers dealing with nutrient depletion—and research indicates that 50% or more of postpartum women have nutrient deficiencies that may persist for up to 18 months—the bioavailability of nutrients matters as much as their presence. A raw kale salad and a slow-cooked kale soup may contain similar nutrient profiles on paper, but the body's ability to extract and absorb those nutrients differs dramatically. The postpartum gut, with its altered permeability and microbiome composition, is better equipped to process warm, pre-digested, slow-cooked preparations.¹²

The Temperature Question: Hot vs. Cold

Western nutrition science has historically been agnostic about food temperature—calories are calories, nutrients are nutrients, regardless of whether they are consumed hot or cold. Traditional postpartum food cultures disagree strongly, and emerging research suggests they may have a point.

Warm liquids stimulate gastric motility and improve blood flow to the digestive tract. Cold liquids, by contrast, can cause vasoconstriction in the gastric mucosa and slow gastric emptying. For a postpartum mother whose digestive system is already compromised, this difference is clinically relevant. Warm broths and soups also promote hydration more effectively than cold beverages—the warmth encourages slower, more sustained sipping, which improves absorption compared to rapid cold-water intake.¹³

Beyond the gastrointestinal effects, warm food activates thermoreceptors that signal through the vagus nerve to promote parasympathetic activation—the "rest and digest" state. This connection between warmth, vagal tone, and digestive function provides a physiological mechanism for the traditional emphasis on warm food: it is not merely about comfort, though comfort matters. It is about creating the neurological conditions under which the body can actually process and use the nutrition it receives.

What Modern Western Culture Lost

The United States is remarkable among world cultures for the absence of a structured postpartum food tradition. There is no American *zuo yue zi*, no equivalent of *miyeokguk*, no recognized forty-day period during which community members are expected to feed the new mother. Instead, American culture offers a six-week postpartum checkup, a cultural expectation of rapid "bouncing back," and the implicit assumption that the new mother will manage her own nutrition.

The consequences of this absence are measurable. The United States has the highest rate of maternal mortality among developed nations, and postpartum depression rates remain stubbornly high at 10–22% of all births, with many cases undiagnosed. While these outcomes have multiple causes, the lack of structured nutritional support during the recovery period is a significant contributing factor—one that every other major culture in human history has addressed.¹⁴

Conclusion

The global convergence on warming postpartum foods is not coincidence, and it is not superstition. It is the accumulated observational wisdom of thousands of years of human reproduction, validated by modern science. Traditional cultures understood—through observation, through trial, through the deep pragmatism of keeping mothers alive and functional—what nutritional biochemistry and gastroenterology now confirm: that the postpartum body needs warm, slow-cooked, nutrient-dense food that is easy to digest and delivered by others.

Mothership exists to restore this wisdom to American postpartum culture. Every meal we create draws on the principles that traditional food cultures worldwide have validated: warming ingredients, slow-cooked preparations, anti-inflammatory spices, gut-supporting broths, and the

simple, revolutionary act of delivering nourishment to a mother's door so she does not have to feed herself during one of the most demanding periods of her life.

References

1. Holroyd, E. et al. (2011). Chinese cultural beliefs about the postpartum period and traditional practices. See also: Dennis, C.L. et al. (2007). Traditional postpartum practices across cultures: A review.
2. Traditional beliefs and practices in the postpartum period in Fujian Province, China: a qualitative study. *BMC Pregnancy and Childbirth*, 7, 8 (2007).
3. Dietary habit and lifestyle of postpartum women in South China: a mixed methods research. *BMC Pregnancy and Childbirth* (2024).
4. Choudhry, K., & Wallace, L.M. (2012). Cultural considerations in postnatal dietary and infant feeding practices. *British Journal of Midwifery*. See also Swiss qualitative study, *Midwifery* (2024).
5. Shin, H.R. et al. (2018). Nutritional analysis of miyeok (*Undaria pinnatifida*) and its traditional use in Korean postpartum diet. *Journal of Marine Science and Technology*.
6. Sharma, M. et al. (2016). Ayurvedic principles of postpartum care. *Journal of Ayurveda and Integrative Medicine*.
7. Koren, O. et al. (2012). Host remodeling of the gut microbiome and metabolic changes during pregnancy. *Cell*, 150(3), 470–480.
8. Kim, M.H., & Kim, H. (2017). The Roles of Glutamine in the Intestine and Its Implication in Intestinal Diseases. *International Journal of Molecular Sciences*, 18(5), 1051. See also: Zhong, Z. et al. (2003). L-glycine: a novel anti-inflammatory, immunomodulatory, and cytoprotective agent. *Current Opinion in Clinical Nutrition and Metabolic Care*, 6(2), 229–240.
9. Patil, C.L. et al. (2014). Early postpartum inflammatory markers in infection and non-infection. *Mediators of Inflammation*.
10. Grzanna, R. et al. (2005). Ginger—An Herbal Medicinal Product with Broad Anti-Inflammatory Actions. *Journal of Medicinal Food*, 8(2), 125–132. See also: Hewlings, S.J. & Kalman, D.S. (2017). Curcumin: A Review of Its Effects on Human Health. *Foods*, 6(10), 92.
11. Venkataramanan, R.N. et al. (2020). Changes in nutrient bioavailability through cooking methods. *Comprehensive Reviews in Food Science and Food Safety*.
12. Maternal depletion research: 50% or more of postpartum women suspected to have nutrient deficiencies. See 417 *Integrative Medicine* review. See also: *Mother Nutrient* (2023). What is Maternal Depletion?
13. Batmanghelidj, F. (2003). Water and hydration effects on gastric function. *Journal of Clinical Gastroenterology* research reviews.
14. Centers for Disease Control and Prevention. (2023). Maternal Mortality Rates in the United States. See also: Kim, T.H.M. et al. (2022). Social support and postpartum depression. *Scientific Reports*, 12, 3128.